THE INFLUENCE OF NIFEDIPINE INDUCTION TO GINGIVAL EPITHELIAL CELL PROLIFERATION (*in vivo* study in rat)

Dewi Agustina¹, AL. Supartinah², Irianiwati³, Suryono⁴

¹Dept. of Oral Medicine, Fac. of Dentistry, University of Gadjah Mada, Yogyakarta, Indonesia.

²Dept. of Pedodontic, Fac. of Dentistry, University of Gadjah Mada, Yogyakarta, Indonesia.

³Dept of Pathology of Anatomy, Fac. of Medicine, University of Gadjah Mada, Yogyakarta, Indonesia.

⁴Dept. of Periodontology, Fac. of Dentistry, University of Gadjah Mada, Yogyakarta, Indonesia.

Dewi Agustina, AL. Supartinah, Irianiwati, Suryono. The influence of nifedipine induction to gingival epithelial cell proferation (*in vivo* study in rat). Journal Dentistry Indonesia 2004; 11 (1): 29-34.

Abstract

It is well known that nifedipine administration in hypertensive patients results in gingival hyperplasia. The aim of this study was to study the pattern of nifedipine-induced gingival hyperplasia, based on morphometric and histological changes as well as on PCNA (Proliferating Cell Nuclear Antigen) expression in the gingival epithelium. In total, 36 male Sprague Dawley rats at the age of 6 – 8 weeks were divided into nine experimental groups and three control groups. Each animal received daily DMSO (dimethyl sulfoxide) *via* oral intubation at a dosage of 0 (for control groups), 15, 30 or 60 mg/ kg (experimental groups) of body weight for 7, 21 or 42 days. After the animals were sacrificed, impression of the lower gingival tissue was taken to measure mesio-distal distance, labio-lingual distance and papilla height. The number of blood vessels and the thickness of gingival epithelium were assessed from hematoxylin and eosin stained sections. Proliferative activity of the epithelial cells was determined by immunohistochemical analysis using PCNA monoclonal antibody. Significant increase in the mesio-distal and labio-lingual distance of the lower gingival tissue was detected morphometrically ($p < 0.05$). There were more blood vessels in the experimental groups than in the control groups, however there was no specific pattern based on the dosage or duration of nifedipine administration. On the other hand, significant differences were found in the gingival epithelial thickness and proliferative activity between the experimental and the control groups. PCNA-positive cells were observed in basal and suprabasal layers, but nearly none in lamina propria.

Introduction

Nifedipine, a calcium channel blocker is widely used for long period of time in the clinical management of hypertension or myocardial disease^{1,2,3,4}. One of side effects of nifedipine utilization was gingival enlargement particularly on labial surface of anterior lower gingiva and of anterior upper gingival^{1,2,5}. However, this enlargement did not occur either on edentulous area or on lingual surface of anterior lower gingival. Clinically, the gingival enlargement was similar to that of verapamil and phenytoin usage⁴, however the occurrence of the gingival enlargement is still controversy⁶. Local factors such as stimulus or dental plaque may exaggerate the occurrence of gingival enlargement⁷. Although drug-induced gingival enlargement is not attributed to tumorigenesis, however some cases identified as squamous cell carcinoma resulting from gingival hyperplasia induced by cyclosporin and phenytoin⁸.

Gingival enlargement prevalence varies from 10–50% because of dilantin utilization⁹, from 13–50% because of phenytoin administration and 15–83% because of nifedipine usage¹⁰. Gingival enlargement is influenced by dosage and duration of drug induction. Dosage of 250 µg/g in diet will lead to gingival enlargement after 20 days administration. However on the fortieth day the graph of enlargement will be flat². Drug dosage is not a determinant factor in the occurrence of gingival enlargement⁶. However nifedipine concentration in the gingival tissue or in its metabolite is more to be an essential variable the gingival

enlargement manifestation¹¹. Drug combination, dosage and duration of administration do not give rise any significantly difference in gingival enlargement¹². In human, gingival enlargement is detected after 1–2 month usage of the drug, in dog it occurs after 7 weeks and in rat happens after 8–9 months¹³.

Mechanism of gingival enlargement induced by nifedipine has been studying intensively. In patient with gingival enlargement, concentration of nifedipine in gingival cervical fluid is higher compared to that in the plasma⁴. In normal condition, fibroblast growth might be controlled and its production is constant because of coordination among transcriptional mechanism, post translation mechanism and its intracellular degradation¹⁴. Collagen synthesis is influenced by hormonal factor and epidermal growth factor (EGF). Phenytoin could stimulate testosterone metabolism and increase 5α dihydrotestosterone level which stimulates fibroblast growth, leads to the proliferation of cellular components. Increased EGF level causes the more responsive cell to EGF and the more stimulation to DNA synthesis². The calcium channel blocker could decrease Ca⁺⁺ influx through cell membrane by decreasing the membrane permeability and by blocking the removal of intracellular Ca⁺⁺ as well. Hence, these give rise to decreasing or blocking of cellular secretion function¹⁰. Nifedipine is assumed to influence directly in gingival fibroblast¹⁵ by forming the connective tissue matrix. Homeostasis disturbances may produce many locus in which medication could interfere in synthesis and degradation of collagen¹⁶. Collagenolytic sign of

inflammatory cells in fibroblast and metaloproteinase are phenomena that depend on calcium. Calcium antagonist might interfere Ca⁺⁺ transportation leading to collagen resorption disturbances. On the other hand, *in vitro* study regarding the effect of nifedipine to normal gingival fibroblast result in decreasing of collagen and protein synthesis⁵. It is also happened on enlargement of gingival fibroblast after induced with verapamil⁴.

Histologically, nifedipine-induced gingival enlargement inclined to be an enlargement with normal composition, it was neither hypertrophy nor hyperplasia². In corium area, hyperplasia of fibroblast was detected however inflammatory cells were not found¹³. Hyperplastic epithelial cells, hyperplastic fibroblast and increasing collagen fibers were detected in connective tissue^{1,17,18}. On the other hand, it was found elongation and proliferation features in rete pegs, increasing the number of capillary vessels and infiltration of inflammatory cells¹⁹. Increasing of collagen fiber indicates that there is an abnormal collagen metabolism and fibroblast enlargement indicates that biosynthesis activity grows up. The gingival enlargement has a relationship with extracellular matrix¹⁸. Cell proliferation could be detected by immunohistochemistry analysis using a specific marker such as PCNA since PCNA is a very important DNA replication protein in signal transduction pathway of keratinocyte division²⁰.

From above description it is essential to study the pattern of gingival hyperplasia based on the morphometric, histological changes and PCNA expression of gingival

epithel resulted from nifedipine induction in different dosage and duration.

Material and Methods

Thirty six male Sprague Dawley rats aged 6 – 8 weeks were used in this study. The gingival tissue of the lower jaw after induced with nifedipine in DMSO at a dosage of 0 mg, 15 mg, 30 mg or 60 mg/kg body weight was observed on the 7th, 21st and 42nd days. After the animal was sacrificed, the lower jaw was taken impression for the measurements of mesio-distal, labio-lingual gingival and papilla height. For the mesio-distal gingival measurement : the widest gingival distance from the right second incisor to the left one was measured; for the labio-lingual gingival measurement : the widest gingival distance from labial to lingual surface of interdental space of first upper incisors was measured; for the height of central papilla was obtained by measuring the distance from the peak of papilla to the most concave surface of margin gingiva of the first incisor. Then, the jaw was fixed in buffered formalin 10% for $\pm$ 24 hours and the labial gingival was excised to be embedded in paraffin. Five micrometer sections were provided and stained with Hematoxylin&Eosin for blood vessels counting and measuring the epithelial thickness. Some other sections were immunohistochemically stained with PCNA to observe the number of proliferative cells.

Results

The measurement of the lower jaw of rat after induced with

nifedipine according to the dosage and duration of induction shown in Table 1.

Table 1. Mean (X) and standard deviation (SD) of the mesio-distal (X₁), labio-lingual (X₂), and the height of central papilla (X₃) of the rat's gingival according to dosage and duration of nifedipine induction

Dosage		Duration					
		7 th day		21 st day		42 nd day	
		X̄	SD	X̄	SD	X̄	SD
15 mg	X ₁	3.767	0.237	3.989	0.040	4.006	0.032
	X ₂	3.342	0.087	3.506	0.122	3.574	0.102
	X ₃	2.160	0.690	2.001	0.252	1.992	0.112
30 mg	X ₁	3.962	0.242	4.332	0.066	4.556	0.096
	X ₂	3.528	0.132	3.751	0.406	3.777	0.294
	X ₃	2.032	0.039	2.041	0.020	2.328	0.050
60 mg	X ₁	4.417	0.136	4.494	0.115	5.103	0.181
	X ₂	3.797	0.055	3.814	0.099	3.828	0.066
	X ₃	2.032	0.237	2.163	0.166	2.339	0.037
Control	X ₁	3.706	0.251	3.999	0.028	4.013	0.007
	X ₂	3.286	0.035	3.529	0.142	3.584	0.080
	X ₃	1.032	0.119	2.099	0.212	1.861	0.220

From the Table 1 was shown a tendency of increasing mesio-distal or labio-lingual measurement, however there was no tendency of increasing papilla height measurement. The result of ANOVA for gingival morphometric was conducted based on the dosage and duration of nifedipine induction as shown in Table 2.

From the Table 2 above

demonstrated that significance difference was found ($p < 0.05$) regarding the increasing mesio-distal and labio-lingual gingival measurement because of nifedipine induction based on the dosage and duration. However no significant difference was performed when the gingival measurement was attributed to the duration of induction and the height of papilla.

Table 2. ANOVA of gingival morphometric according to the dosage and duration of nifedipine induction

		F	p
Inter-dosage	x ₁	44.722	0.000
	x ₂	20.547	0.000
	x ₃	12.107	0.000
Inter-duration	x ₁	17.232	0.000
	x ₂	13.338	0.000
	x ₃	2.598	0.078
Dosage-duration	x ₁	49.040	0.000
	x ₂	12.612	0.000
	x ₃	9.609	0.000

Fig. 1 : Rat's gingival epithel with high rete pegs (arrow) (H&E, X400).

Fig. 2 : Immunohistochemistry staining using PCNA monoclonal antibody in rat's gingival epithel (X1000). Positive stainings were shown in nuclei appeared as dark brown stainings (arrows) especially in basal and parabasal layers accompanied with loose connective tissue.

Table 3. Mean and standard deviation of the number of blood vessel and epithelial thickness measurement based on the dosage and duration of nifedipine induction.

Dosage (mg) / Duration (day)	Blood vessel	Epithelial thickness (mm)
15/7	1.440 ± 0.191	0.124 ± 0.038
15/21	2.107 ± 0.387	0.146 ± 0.016
15/42	1.217 ± 0.510	0.129 ± 0.005
30/7	1.993 ± 0.577	0.143 ± 0.020
30/21	1.330 ± 0.33	0.129 ± 0.045
30/42	1.773 ± 0.836	0.159 ± 0.009
60/7	1.443 ± 0.709	0.159 ± 0.008
60/21	1.773 ± 0.196	0.147 ± 0.013
60/42	1.550 ± 0.191	0.183 ± 0.023
Control	1.663 ± 0.57	0.110 ± 0.011

The result of blood vessels enumeration and epithelial thickness measurement are shown in Table 3.

From the table above shown that the number of blood vessel in experimental group was higher compared to the control group. However the increasing of that did not describe any particular pattern either according to the dosage or duration of nifedipine induction. From the ANOVA demonstrated no significant difference ($p > 0.05$).

It was clear that the epithel of the experimental group was thicker than that of the control group and it was also demonstrated in ANOVA based on the dosage and duration of nifedipine induction ($p < 0.05$). The result of ANOVA from the enumeration of blood vessel and the epithelial thickness appear in Table 4.

Proliferative activity of epithelial cell was determined by counting the number of positive staining cell analysed by immunohistochemistry using PCNA monoclonal antibody from 100 gingival epithelial cells for 1000X magnification and it was repeated for three times per section. The result are demonstrated in Table 5.

From the table above seen that the higher dosage and the longer of nifedipine induction gave the more positive PCNA-stained cells. The most positive cell was found at the dosage of 60 mg for 7 days. By using ANOVA the proliferative index of epithelial cell are given in the table below.

From the table above shown that among the dosage, duration and interaction of both gave significant difference regarding the proliferative activity of gingival epithelial cells.

Table 4. ANOVA of the enumeration of blood vessel and the epithelial thickness according to the dosage and duration of nifedipine induction

	F	P
Inter - dosage		
- Blood vessel	0.070	0.975
- Epithelial thickness	6.752	0.005
Inter - duration		
- Blood vessel	0.207	0.890
- Epithelial thickness	4.197	0.026
Dosage - duration		
- Blood vessel		
- Epithelial thickness	0.579	0.794
	1.764	0.164

Table 5. Mean and standard deviation of proliferative activity of gingival epithelial cells using PCNA - staining according to the dosage and duration of nifedipine induction

Dosage (mg/day)	Mean $\pm$ SD
15/7	41.67 $\pm$ 8.50
15/21	23.00 $\pm$ 1.73
15/42	33.67 $\pm$ 3.51
30/7	47.67 $\pm$ 6.65
30/21	57.33 $\pm$ 4.93
30/42	57.00 $\pm$ 10.58
60/7	59.77 $\pm$ 3.42
60/21	54.00 $\pm$ 2.00
60/42	54.67 $\pm$ 3.78
control	44.67 $\pm$ 2.88

Table 6. ANOVA of epithelial cell proliferative index according to the dosage and duration of nifedipine induction

	F	p
Inter - dosage	34.421	0.000
Inter - duration	3.750	0.036
Dosage - duration	3.053	0.030

Discussion

From the result of this study, it was apparent that morphological gingival size significantly increased especially for mesio-distal and labio-ligual gingival distances. Histologically, the connective tissue became loose, proliferation was dominantly performed in basal and suprabasal cells as detected immunohistologically. The number of blood vessel did not significantly increase yet, but the epithelial thickness increased significantly.

If there is an excessive or continuous stimulation, a cell will adjust to maintain homeostasis by increasing metabolic activity characterized by accumulation of intra and extra-cellular matrix that clinically manifested as hyperplasia, hypertrophy, atrophy or metaplasia²¹.

Gingival hyperplasia because of nifedipine induction is caused by involvement of alteration of Na^+ and Ca^{++} flow, of folic acid, of collagenase activation and local factor such as bacterial plaque. Nifedipine results in decreasing of Na^+ flow either direct or indirect depends on Na^+ and Ca^{++} exchange. This causes decreasing of folic acid that results in intracellular folic acid deficiency. In turn, production of *trypsin-like collagenase activator* will be limited, so stock of collagenase will be limited, as well¹⁷.

Fibroblast and endothelial cell of vascular tissue are classified as stabil cells because of their low replication level, however these cells might replicate rapidly if appropriate stimulation exists²¹. The statement above could be used as a reason why there was no significant difference of the number of blood vessels between the experimental and the control

groups. Nifedipine action in bothering the Ca^{++} influx into cell was assumed as a stimulation to fibroblast in extracellular matrix formation. Because of limited collagenase product it results in enlargement of tissue morphologically¹⁸.

Oral epithelial cell denotes labil cell and has high proliferative capacity²¹. Nifedipine induction will stimulate basal and parabasal cells of gingival epithel to become more active in proliferation that resulted in increasing gingival epithelial thickness. It has been reported that *c-myc* expression could be detected in nucleus of basal and parabasal cells of the nifedipine/phenytoin-induced gingival hyperplasia²². It is reasonable since *myc* is the first nuclear gene involved in the kinase cascade of oral epithelial cell division²⁰.

From the histological finding in this study demonstrated loose connective tissue, probably as a result of Ca^{++} influx disturbances that is clinically seen as tissue's oedema. On the other hand, indeed, there was increasing number of blood vessels in the experimental group, however it was clear that no specific pattern appeared according to the dosage or duration of nifedipine induction. This result suggested that no obvious inflammatory process involved in this nifedipine-induced gingival enlargement. The latter result supported a finding that histologically nifedipine induction inclined to be an enlargement with normal composition neither hypertrophic or hyperplastic process². This tenet also confirmed a result of study that in nifedipine induced tissue did not find any inflammatory cell¹³. Increasing number of the new blood vessels as a result of new capillary opening that in

The latter result supported a finding that histologically nifedipine induction inclined to be an enlargement with normal composition neither hypertrophic or hyperplastic process². This tenet also confirmed a result of study that in nifedipine-induced tissue did not find any inflammatory cell¹³. Increasing number of the new blood vessels as a result of new capillary opening that in normal condition those capillaries are closed²¹. The opening of the capillary is considered as the additional capillary as the result of rete pegs elongation since there was an increasing metabolic activity because of nifedipine induction.

Conclusions

Nifedipine induction resulted in increasing gingival size particularly on labial surface area of the anterior lower jaw. On the other hand, cellular proliferative rate of gingival epithel increased significantly especially in basal and suprabasal layer. However, the enlargement neither involved inflammatory or hypertrophic/hyperplastic processes, it tended to be an enlargement with normal composition.

References

1. Nishikawa S, Tada H, Hamasaki A, Kasahara S, Kido J, Nagata T, Ishida H, Wakano, Y. Nifedipine-induced gingival hyperplasia: A clinical and *in vitro* study. *J Periodontol* 1991. 62 : 30-35.
2. Nishikawa S, Nagata T, Morisaki I, Oka T, Ishida H. Pathogenesis of drug-induced gingival overgrowth. A review of studies in the rat model. *J Periodontol* 1996. 67 : 463-71.
3. Barber MT, Savage NW and Seymour GJ. The effect of cyclosporin and lipopoly-saccharide on fibroblast: implications for cyclosporin-induced gingival overgrowth. *J Periodontol* 1992. 63 : 397-404.
4. Fu E, Nieh S, Hsiao CT, Hsieh YD and Wikesyo UME. Nifedipine-induced gingival overgrowth in rats: brief review and experimental study. *J Periodontol* 1998. 69 : 765-71.
5. Sato T, Oikarinen KS, Oikarinen AI. Effect of phenytoin and nifedipine on collagen gene expression in human gingival fibroblasts. *J Oral Pathol Med* 1990. 19 : 404-7.
6. Conard GJ, Jeffay H, Boshes L and Steinberg, AD. Levels of 5,5-Diphenylhydantoin and its major metabolite in human serum, saliva and hyperplastic gingival. *J Dent Res* 1974. 53 : 1323-29.
7. Morisaki I, Kato, Loyola-Rodriguez JP, Nagata T, Ishida H. Nifedipine-induced gingival overgrowth in the presence or absence of gingival inflammation in rats. *J Periodont Res* 1993. 29 : 396-403.
8. Saito K, Mori S, Tanda N, Sakamoto S. Expression of p53 protein and Ki-67 antigen in gingival hiperplasia induced by nifedipine and phenytoin. *J Periodontol* 1999. 70 : 581-6.
9. Seymour RA, Smith DG, Rogers SR. The comparative effects of azathioprine and cyclosporine on some gingival health parameters of renal transplant patients. *J Clin Periodontol* 1987. 14 : 610-3.
10. Barclay S, Thomaso, JM, Idle JR and Seymour RA. The incidence and severity of nifedipine-induced gingival overgrowth. *Clin Periodontol*. 1992. 19 : 311-4.
11. Cimasoni G. Crevicular fluid updated. In: Myers HM (ed) : *Monographs in oral science*, vol. 12. Base: S.Karger. 1983: 1-52.
12. Tavassoli S, Yamalik N, Caglayan F, Caglayan G and Eratalay K. The clinical effects on nifedipine on periodontal status. *J Periodontol* 1998. 69 : 108-12.
13. Nyska A, Waner T, Pirak M, Galiano A and Zlatogorski A. Gingival hyperplasia in rats induced by oxodipine - a calcium channel blocker. *J Periodont Res* 1990. 25 : 62-8.
14. Narayanan AS, Meyers DF and Page RC. Regulation of collagen production in fibroblast cultured from normal and phenytoin-induced hyperplastic human gingiva *J Periodont Res* 1988. 23 : 118-21.
15. van der Wall E, Tuinzing DB and Hes J. Gingival hyperplasia induced by nifedipine, an arterial vasodilating drug. *Oral Surg, Oral Med, Oral Pathol* 1985. 60 : 38-40.
16. Uitto J, Bauer EA, Santa Cruz D, Holtmann B and Eissen AZ. Decreased collagenase production by regional fibroblasts cultured from tissue nevi of the collagen type. *J Inv Dermatol* 1982. 78 : 139-40.
17. Brown RS, Beaver WT and Bottomley. On the mechanism of drug-induced gingival hyperplasia. *J. Oral Pathol Med* 1991. 20 : 201-9.
18. Tipton DA, Fry HR and Dabbous MKH. Altered collagen metabolism in nifedipine-induced gingival overgrowth. *J Periodont Res* 1994. 29 : 401-9.
19. Romanos GE, Strub JR and Bernimoulin JP. Immunohistochemical Distribution of Extracellular Matrix Proteins as a Diagnostic Parameter in Healthy and Diseased Gingiva. *J Periodontol* 1993. 64 : 110-9.
20. Sugerman PB and Savage NW. Current concepts in oral cancer. *Australian Dental Journal* 44 : 147 - 156.
21. Cotran RS, Kumar V and Robbins SL. *Pathologic Basis of Disease*. Philadelphia WB Saunders Co 1994. 36-7, 44-9, 469-71.
22. Saito K, Mori S, Tanda N and Sakamoto S. Immunolocalization of c-Myc and bcl-2 protooncogene products in gingival hiperplasia induced by nifedipine and phenytoin. *J Periodontol* 2000. 71 : 44-9.

Acknowledgement

The authors wish to thank to Badan Penelitian dan Pengembangan Kesehatan (Medical Research and Development Agency), Department of Health, Republic of Indonesia which has provided the funding and to Lembaga Biologi Molekuler Eijkman (Molecular Biological Institution of Eijkman) which has given a chance for this study undertaken.